

Western Trauma Association critical decisions in trauma: Management of the open abdomen after damage control surgery

Jack Sava, MD, Hasan B. Alam, MD, Gary Vercrucyse, MD, Matthew Martin, MD, Carlos V. R. Brown, MD, Karen Brasel, MD, MPH, Ernest E. Moore, Jr, MD, David Ciesla, MD, Kenji Inaba, MD, and for the Western Trauma Association Critical Decisions in Trauma Committee, Washington, District of Columbia

Damage control laparotomy has become the standard of care in the management of severely injured patients. The popularization of this approach has raised many important issues regarding technique, timing, and closure of the open abdomen (OA). Substantial practice variation exists, and improvised or surgeon-constructed dressings now compete with commercially available solutions. Choices in managing patient with an OA may have a profound impact on the outcome of the abdominal wall, but— even more importantly—may affect mortality as well.¹

This algorithm synthesizes published evidence with the expert opinion of members of the Western Trauma Association (WTA) members, presenting a systematic approach for the management of the OA after damage control laparotomy. These recommendations are created using an iterative process (Fig. 1) beginning with a detailed review of the literature by the members of the WTA Critical Decisions in Trauma Committee (“Algorithm Committee”). Initial draft recommendations are refined by Algorithm Committee members, then vetted at the WTA Annual Meeting, before final review by the Committee.

A: GOALS

The goal of this algorithm is to provide an easily understandable and practical protocol that can be used by surgeons and other clinicians to better manage patients with severe OAs. Where evidence is mixed or insufficient, the algorithm uses expert opinion to guide trauma providers towards safe, widely-accepted clinical decisions and strategies.

B: BURDEN OF DISEASE

While use of OA techniques has increased in recent decades, the number of patients undergoing temporary abdominal closure is not known. Up to 25% of patients undergoing

emergent laparotomy for injury may be unsuitable for primary closure at the index operation,² although some authors have reported that consensus does not exist for indications, and OA techniques should be applied more sparingly.^{3–5} Patients with OA may be expected to incur a rate of enterocutaneous fistula of approximately 8%.⁶ Patients who cannot be closed primarily must be discharged with a planned hernia, leading to an additional subsequent elective operation and hospitalization for hernia repair.

C: EXISTING GUIDELINES

Several reviews summarize available evidence on the spectrum of issues relevant to OA management.⁷ The World Society for Emergency Surgery has published an excellent, exhaustive evidence-based guideline with graded recommendations.⁸ Coccolini et al.⁹ have published a detailed and well-illustrated book on the topic. The goal of the WTA algorithm is to provide a practical, simple algorithm that synthesizes available evidence as well as expert opinion of members of the WTA and the algorithms committee (Fig. 2).

E: OA MANAGEMENT IN EMERGENCY GENERAL SURGERY

While damage control techniques have been more widely studied in trauma, the technique has gained broad usage in a variety of emergency surgical settings.^{3,8,10,11} Indications in this context may parallel those used in trauma operations, but may more commonly focus on the need to limit exposure to general anesthesia and operative fluid loss in critically ill patients, especially those with little physiologic reserve. Previous WTA algorithms have acknowledged the use of damage control in nontrauma settings¹² but this algorithm focuses on OA management in injured patients.

F: DEFINITIONS, CONCEPTS AND THEORETICAL RATIONALES

- a. Open abdomen: For this manuscript, this term applies when damage control laparotomy is performed without primary fascial closure at the initial operation, with a plan for interval, staged relaparotomy.

Submitted: March 2, 2019, Revised: May 8, 2019, Accepted: May 9, 2019, Published online: June 12, 2019.

From the MedStar Washington Hospital Center (J.S., H.B.A., G.V., M.M., C.V.R.B., K.B., E.E.M., D.C., K.I.), Washington, District of Columbia.

This algorithm was presented for discussion at the 48th Annual Meeting of the Western Trauma Association, Feb 25–March 2, 2018, Whistler-Blackcomb, British Columbia, Canada.

Address for reprints: Jack Sava, MD, MedStar Washington Hospital Center, Room 4B39, 110 Irving St., NW Washington, DC 20010; email: jack.a.sava@medstar.net.

DOI: 10.1097/TA.0000000000002389




Figure 1. Process for creation of WTA critical decision algorithm.

- b. Packing: temporary use of nonabsorbable, nonimplantable gauze to promote hemostasis
- c. Primary fascial closure: full length fascia-to-fascia approximation, with or without reinforcement.
- d. Progressive abdominal closure: A strategy of serial, staged operations aimed at progressively approximating fascial edges to facilitate primary fascial closure

G. MANAGEMENT

(1) Damage Control/OA may be indicated during laparotomy for trauma if any of the following circumstances are noted:

- (a) Physiologic indicators suggest that operation should be abbreviated. Persistent hypotension, acidosis, core temperature less than 34°C, and coagulopathy may signal impending death, and should prompt a damage control approach.¹³⁻¹⁷ Historically, pH has been used to describe acidemia thresholds for damage control (pH < 7.2), but more accurate measures of metabolic acid load such as lactate and base deficit are widely used and well validated.^{18,19} Coagulopathy has traditionally been diagnosed via clinical observation of the surgical field, or by standard laboratory values (PTT/INR/platelet count/fibrinogen). However, recent work suggests that viscoelastic assays (thromboelastography and thromboelastometry may be important adjuncts in the assessment of clinical




Figure 2. WTA algorithm: management of OA after damage control.

coagulopathy.²⁰⁻²² Other point-of-care tools (e.g., impedance aggregometry) may have an evolving role,²³ but are not currently in common clinical use.

Surgical bleeding must be controlled prior to ending the operation, but endless futile searches for sources of diffuse

coagulopathic hemorrhage are to be avoided. Peritoneal lavage for rewarming can be effective in reversing core hypothermia, but is disruptive to packing, and its effect on outcome is not known. Physiologic trends—in addition to initial values—can help guide the decision for damage control, and with appropriate resuscitation and prompt hemorrhage control some patients

may avoid damage control despite significant early physiologic derangement.

(b) Abdomen cannot be closed without abdominal hypertension.¹⁶ Excessive fascial tension will often be palpably evident to the surgeon at the time of closure. Bladder pressure may be measured, but is cumbersome in this setting. Peak airway pressure is often measured during fascial approximation, though some data suggests poor correlation with intra-abdominal pressure.²⁴ Exact cutoff values permitting closure are not known, and judgment and experience should guide the decision to proceed.

(c) The patient is at increased risk for postoperative abdominal compartment syndrome. Even if excessive fascial tension is not present at the time of planned closure, risk of later development of intraabdominal hypertension (IAH) may be unacceptably high. Early IAH may impact important outcomes including survival.²⁴ Numerous potential risk factors have been described among injured patients as well as other critically ill populations. While there is no universal consensus, the following are likely contributors to abdominal hypertension after injury:

1. Severe shock/resuscitation (hypotension, serum lactate >5, blood loss >4 L, transfusion >10 units packed red cells)^{7,25–27}
2. Injuries requiring packing^{28,29}
3. Obvious visceral swelling
4. Abdominal wall tissue loss
5. Obesity²⁶

(d) Additional delayed definitive abdominal procedures (e.g., bowel anastomosis) are required. Even in the absence of abdominal hypertension, fascial closure should be deferred if relaparotomy for further organ repair/debridement is planned. Repeated closing and reopening of fascia causes unnecessary tissue damage.

(e) Nonabsorbable intraabdominal packing is used, or temporary devices (e.g., vascular shunts) are left in the abdomen. If the decision is made to manage hemorrhage with packing, a temporary abdominal closure should be used which allows for abdominal swelling without excessive pressure increases. A balance exists between using packs to create local positive pressure/tamponade, while avoiding compression of major vessels and creation of abdominal hypertension.

(f) A second look operation is needed (e.g., marginal bowel viability). Planned reevaluation of marginally perfused bowel or other organs may be appropriate if clear demarcation is not visible, or when abdominal viscera are globally hypoperfused due to shock, making accurate evaluation difficult.

If an indication for damage control does not exist, fascia should be closed primarily at the first operation to avoid the

deleterious consequences of the OA. Many emergent trauma operations (e.g., uncomplicated splenectomy for severe injury with transient hypotension) will not require damage control. Physiology should be continuously reassessed during surgery, and consideration given to primary closure if patient condition improves over the course of the operation. Patients without packing who are easily closed at the second operation may benefit from review for appropriateness of damage control, as part of an institutional continuous quality improvement program.⁵

(2) Choice of dressing and attention to detail in critical care management have significant impact on outcomes

(a) The initial dressing may be a commercial product (e.g., Abthera, KCI), or may be fashioned by the surgeon from available surgical supplies.^{30–34} In either event, it should be watertight, and should employ negative pressure to drain abdominal fluid and allow for its measurement. The dressing should be lax enough to allow for abdominal swelling without abdominal hypertension. The layer adjacent to viscera should be smooth and nonadherent. Additional closed suction drains may be placed as part of specific injury management. Primary fascial closure should not be performed in the damage control setting. Temporary closure of skin (e.g., with towel clamps) may put the patient at risk for subsequent abdominal hypertension.³⁵

(b) In the perioperative period, principles of damage control resuscitation should be applied. Fluid balance should be measured carefully, including abdominal losses. Excessive crystalloid should be avoided.³⁶ Diuresis may be useful in selected cases, but routine aggressive diuresis has no demonstrated efficacy.³⁷ Balanced blood products should be used to correct anemia and coagulopathy, guided by predetermined ratios or laboratory tests/thromboelastography.²⁰ Intraabdominal pressure should be measured, as IAH or abdominal compartment syndrome may develop even with an OA.

(c) Though evidence is limited, Direct Peritoneal Resuscitation—in which hypertonic peritoneal dialysis solution is instilled into the abdominal cavity in the immediate postoperative period—may lead to increased rates of primary fascial closure, as well as improving other physiologic outcomes. Suggested mechanisms include visceral vasodilatation, reduced organ edema, and decreased cytokine levels.³⁸

(d) There is no consensus on prophylactic antibiotic use in OA patients. When used, antibiotics should be appropriate for injuries, and limited in duration.

(e) Appropriately selected patients who pass spontaneous breathing trials may be extubated during OA management.³⁹ Extubation may be deferred if respiratory mechanics are poor, if there is persistent metabolic acidosis, if intervals between repeat operations are short (<24 h), or if a difficult airway is present.

(f) Early enteral nutrition should be provided whenever possible. Damage control patients are hypermetabolic and require careful attention to nutritional needs.⁴⁰ Historically, many surgeons were concerned that gut function might be ineffective when viscera were exposed, or that early feeding might worsen ileus, which might exacerbate loss of domain. However, subsequent experience and studies have shown that early enteral feeding is feasible.^{34,41,42}

(3) Evidence of good response may include stabilization of vital signs, resolution of metabolic acidosis, improvement of end organ function (mental status, urine output, skin perfusion), or other clinical/laboratory markers. Ongoing, uncontrolled hemorrhage may manifest through high volume sanguineous dressing/drain output, bleeding at external wound site, or falling hemoglobin levels. Continued or accelerated bleeding should prompt surgical reexploration to search for surgical correctable bleeding sites. In cases of refractory coagulopathic bleeding, reexploration is often fruitless.

(4) Reoperation should take place as soon as possible after adequate resuscitation and correction of metabolic abnormalities. Ideally, packs are removed within 24 hours, though timing of removal may depend on the indication for packing and the degree of soiling. Excessive delay in reoperation should especially be avoided in patients with proximal bowel in discontinuity, retained packs/sponges, or intravascular shunts.

(5) Patients with prolonged OA may be at added risk for complications including colonic anastomotic leak,⁴³ surgical site infection, fluid loss/metabolic derangement, and failure of primary fascial closure.⁴⁴ Colonic anastomotic leak rates have been found to increase with duration of OA, and when more than one relaparotomy is required,⁴⁵ leading many surgeons to believe that colostomy should be considered if the abdomen is likely to be left open longer than 36 hours, or if multiple relaparotomies are needed. Anastomosis beyond first relaparotomy has been successfully offered by some authors in carefully selected patients.⁴⁶

(6) Progressive abdominal closure—with eventual primary fascial approximation—may be facilitated by:

- (a) continued damage control resuscitation, including judicious management of fluid balance, avoiding excessive crystalloid use;
- (b) application of negative pressure and continuous tension to abdominal wall, using handmade or commercially available (e.g., Wittman patch, Abthera) devices^{47,48,49}. Continuous negative pressure and retention sutures both provide fascial tension, and may be used together^{50,51};
- (c) Progressive sequential closure of fascial edges starting at each end of fascial incision.⁵⁰ Typically, patients return to OR every 2 days for further closure.
- (d) Periodic readjustment of tensioning device, either at bedside or in operating room.

(7) Progressive primary closure should be abandoned when there is lack of progress upon attempts at retensioning, or when duration of OA becomes prolonged (e.g., 7–10 days). Further attempts at reexploration and closure are associated with diminishing likelihood of fascial closure⁵² may put the patient at risk for bowel injury, fistula development, or septic complications^{53,54}

Alternative abdominal closure techniques include:

- (a) Placement of permanent, bridging prosthetic or bioprosthetic mesh. A full discussion of mesh choices is beyond the scope of this document, but simple first generation, wide-pore polypropylene mesh (e.g., Marlex, Prolene) is prone to erosion into intestine, and should not be used directly adjacent to bowel. However, a wide and evolving variety of composite meshes are available with nonerosive layers for contact with bowel.⁵⁵ Bioprosthetics are less prone to erosion, but are associated with high eventration rates when used as a bridge.⁵⁶
- (b) Placement of absorbable mesh such as Vicryl (Polyglactin 910), with or without skin closure or interval split thickness skin grafting
- (c) Skin only closure. Running or interrupted technique may be used. Local skin complications such as necrosis at suture entry sites are common, but rarely catastrophic.
- (d) Complex free/rotational flap closure (typically in cases of massive loss of abdominal wall soft tissue). Choice of flap depends on size and other injuries at potential donor sites.
- (e) Components separation during initial admission has been described⁵⁷ but is more commonly avoided, allowing for use in later elective abdominal wall reconstruction.

DISCLOSURE

The authors declare no funding or conflicts of interest. The opinions expressed in this article are those of the authors and do not reflect the official policy of the affiliated institutions or any government agencies.

REFERENCES

1. Cheatham ML, Safcsak K. Is the evolving management of intra-abdominal hypertension and abdominal compartment syndrome improving survival? *Crit Care Med*. 2010;38(2):402–407.
2. Miller RS, Morris JA Jr., Diaz JJ Jr., Herring MB, May AK. Complications after 344 damage-control open celiotomies. *J Trauma*. 2005;59(6):1365–1371.
3. Khan A, Hsee L, Mathur S, Civil I. Damage-control laparotomy in nontrauma patients: Review of indications and outcomes. *J Trauma Acute Care Surg*. 2013;75(3):365.
4. Higa G, Friese R, O'Keefe T, Wynne J, Bowlby P, Ziembra M, Latifi R, Kulvatunyou N, Rhee P. Damage control laparotomy: a vital tool once overused. *J Trauma*. 2010;69(1):53–59.
5. Harvin JA, Sharpe JP, Croce MA, et al. Better understanding the utilization of damage control laparotomy: a multi-institutional quality improvement project. *J Trauma Acute Care Surg*. 2019; Apr 18 [cited 2019 Apr 26]; Publish Ahead of Print. Available from: https://journals.lww.com/jtrauma/Abstract/publishahead/Better_Understanding_the_Utilization_of_Damage.98336.aspx#pdf-link.

6. Fischer PE, Fabian TC, Magnotti LJ, Schroepfel TJ, Bee TK, Maish GO, Savage SA, Laing AE, Barker AB, Croce MA. A ten-year review of enterocutaneous fistulas after laparotomy for trauma. *J Trauma—Inj Infect Crit Care*. 2009;67(5):924–928.
7. Regner JL, Kobayashi L, Coimbra R. Surgical strategies for management of the open abdomen. *World J Surg*. 2012;36(3):497–510.
8. Coccolini F, Roberts D, Ansaloni L, et al. The open abdomen in trauma and non-trauma patients: WSES guidelines. *World J Emerg Surg*. 2018; 13(1):7.
9. Coccolini F, Ivatury R, Sugrue M, Ansaloni L. *Open Abdomen: A Comprehensive Practical Manual*. Springer; 2018: 281 p.
10. Bruns BR, Ahmad SA, O'Meara L, Tesoriero R, Lauerman M, Klyushnenkova E, Kozar R, Scalea TM, Diaz JJ. Nontrauma open abdomens: a prospective observational study. *J Trauma Acute Care Surg*. 2016;80(4):631–636.
11. Waibel BH, Rotondo MF. Damage control for intra-abdominal sepsis. *Surg Clin North Am*. 2012;92(2):243–257.
12. Moore FA, Moore EE, Burlew CC, Coimbra R, McIntyre RC, Davis JW, Sperry J, Biffl WL. Western Trauma Association critical decisions in trauma: management of complicated diverticulitis. *J Trauma Acute Care Surg*. 2012; 73(6):1365–1371.
13. Sagraves SG, Toschlog EA, Rotondo MF. Damage control surgery—the intensivist's role. *J Intensive Care Med*. 2006;21(1):5–16.
14. Moore EE, Burch JM, Franciose RJ, Offner PJ, Biffl WL. Staged physiologic restoration and damage control surgery. *World J Surg*. 1998;22(12): 1184–1190; discussion 1190–1191.
15. Jurkovich GJ, Greiser WB, Luteran A, Curreri PW. Hypothermia in trauma victims: an ominous predictor of survival. *J Trauma*. 1987;27(9):1019–1024.
16. Roberts DJ, Bobrovitz N, Zygun DA, et al. Indications for use of damage control surgery in civilian trauma patients. A content analysis and expert appropriateness rating study. *Ann Surg*. 2016;263(5):1018–1027.
17. Roberts DJ, Bobrovitz N, Zygun DA, Ball CG, Kirkpatrick AW, Faris PD, Stelfox HT. Indications for use of damage control surgery and damage control interventions in civilian trauma patients: a scoping review. *J Trauma Acute Care Surg*. 2015;78:1187–1196.
18. Davis JW, Kaups KL, Parks SN. Base deficit is superior to pH in evaluating clearance of acidosis after traumatic shock. *J Trauma*. 1998;44(1):114–118.
19. Callaway DW, Shapiro NI, Donnino MW, Baker C, Rosen CL. Serum lactate and base deficit as predictors of mortality in normotensive elderly blunt trauma patients. *J Trauma*. 2009;66(4):1040–1044.
20. Da Luz LT, Nascimento B, Shankarakutty AK, Rizoli S, Adhikari NK. Effect of thromboelastography (TEG®) and rotational thromboelastometry (ROTEM®) on diagnosis of coagulopathy, transfusion guidance and mortality in trauma: descriptive systematic review. *Crit Care*. 2014;18(5):518 [cited 2018 Oct 17];18(5). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4206701/>.
21. Gonzalez E, Moore EE, Moore HB, et al. Goal-directed hemostatic resuscitation of trauma-induced coagulopathy: a pragmatic randomized clinical trial comparing a viscoelastic assay to conventional coagulation assays. *Ann Surg*. 2016;263(6):1051–1059.
22. Hanke A, Horstmann A, Wilhelm A. Point-of-care monitoring for the management of trauma-induced bleeding. *Curr Opin Anaesthesiol*. 2017;30(2): 250–256.
23. Connelly CR, Yonge JD, McCully SP, et al. Assessment of three point-of-care platelet function assays in adult trauma patients. *J Surg Res*. 2017; 212:260–269.
24. Ivatury R, Porter J, Simon R, Islam S, John R, Stahl W. Intra-abdominal hypertension after life-threatening penetrating abdominal trauma: prophylaxis, incidence, and clinical relevance to gastric mucosal pH and abdominal compartment syndrome. *J Trauma Acute Care Surg*. 1998;44(6):1016.
25. Holodinsky JK, Roberts DJ, Ball CG, Blaser AR, Starkopf J, Zygun DA, Stelfox HT, Malbrain ML, Jaeschke RC, Kirkpatrick AW. Risk factors for intra-abdominal hypertension and abdominal compartment syndrome among adult intensive care unit patients: a systematic review and meta-analysis. *Crit Care*. 2013;17(5):R249.
26. Kim IB, Prowle J, Baldwin I, Bellomo R. Incidence, risk factors and outcome associations of intra-abdominal hypertension in critically ill patients. *Anaesth Intensive Care*. 2012;40(1):79–89.
27. Vidal M, Weisser J, Gonzalez F, Toro M, Loudet C, Balasini C, Canales H, Reina R, Estenssoro E. Incidence and clinical effects of intra-abdominal hypertension in critically ill patients. *Crit Care Med*. 2008;36(6): 1823–1831.
28. Meldrum DR, Moore FA, Moore EE, Franciose RJ, Sauaia A, Burch JM. Prospective characterization and selective management of the abdominal compartment syndrome. *Am J Surg*. 1997;174(6):667–673.
29. Meldrum DR, Moore FA, Moore EE, Haanel JB, Cosgriff N, Burch JM. Cardiopulmonary hazards of perihepatic packing for major liver injuries. *Am J Surg*. 1995;170(6):537–542.
30. Campbell AM, Kuhn WP, Barker P. Vacuum-assisted closure of the open abdomen in a resource-limited setting. *S Afr J Surg*. 2010;48:114–115.
31. Navsaria PH, Bunting M, Omshoro-Jones J, Nicol AJ, Kahn D. Temporary closure of open abdominal wounds by the modified sandwich–vacuum pack technique. *BJS*. 2003;90(6):718–722.
32. Brock WB, Barker DE, Burns RP. Temporary closure of open abdominal wounds: the vacuum pack. *Am Surg*. 1995;61(1):30–35.
33. Foy HM, Nathens AB, Maser B, Mathur S, Jurkovich GJ. Reinforced silicone elastomer sheeting, an improved method of temporary abdominal closure in damage control laparotomy. *Am J Surg*. 2003;185(5):498–501.
34. Barker DE, Kaufman HJ, Smith LA, Ciraulo DL, Richart CL, Burns RP. Vacuum pack technique of temporary abdominal closure: a 7-year experience with 112 patients. *J Trauma Acute Care Surg*. 2000;48(2):201.
35. Raeburn CD, Moore EE, Biffl WL, Johnson JL, Meldrum DR, Offner PJ, Franciose RJ, Burch JM. The abdominal compartment syndrome is a morbid complication of postinjury damage control surgery. *Am J Surg*. 2001;182(6): 542–546.
36. Ghneim MH, Regner JL, Jupiter DC, Kang F, Bonner GL, Bready MS, Frazee R, Ciceri D, Davis ML. Goal directed fluid resuscitation decreases time for lactate clearance and facilitates early fascial closure in damage control surgery. *Am J Surg*. 2013;206:995–999.
37. Webb LH, Patel MB, Dortch MJ, Miller RS, Gunter OL, Collier BR. Use of a furosemide drip does not improve earlier primary fascial closure in the open abdomen. *J Emerg Trauma Shock*. 2012;5(2):126–130.
38. Weaver JL, Smith JW. Direct peritoneal resuscitation: a review. *Int J Surg*. 2016;33:237–241.
39. Sujka J, Safcsak K, Cheatham M, Ibrahim J. Trauma patients with an open abdomen following damage control laparotomy can be extubated prior to abdominal closure. *World J Surg*. 2018;42(10):3210–3214.
40. Cheatham ML, Safcsak K, Brzezinski SJ, Lube MW. Nitrogen balance, protein loss, and the open abdomen*. *Crit Care Med*. 2007;35(1):127.
41. Burlew CC, Moore EE, Cuschieri J, et al. Who should we feed? A Western Trauma Association multi-institutional study of enteral nutrition in the open abdomen after injury. *J Trauma Acute Care Surg*. 2012;73(6):1380.
42. Byrnes MC, Reicks P, Irwin E. Early enteral nutrition can be successfully implemented in trauma patients with an “open abdomen”. *Am J Surg*. 2010; 199:359–362; discussion 363.
43. Chamieh J, Prakash P, Symons WJ. Management of destructive colon injuries after damage control surgery. *Clin Colon Rectal Surg*. 2018;31(1): 36–40.
44. Swaroop M, Williams M, Greene WR, Sava J, Park K, Wang D. Multiple laparotomies are a predictor of fascial dehiscence in the setting of severe trauma. *Am Surg*. 2005;71(5):402–405.
45. Anjaria D, Ullmann T, Lavery R, Livingston DH. Management of colonic injuries in the setting of damage-control laparotomy: one shot to get it right. *J Trauma Acute Care Surg*. 2014;76:594–600.
46. Ordoñez CA, Pino LF, Badiel M, Sánchez AI, Loaiza J, Ballestas L, Puyana JC. Safety of performing a delayed anastomosis during damage control laparotomy in patients with destructive colon injuries. *J Trauma Inj Infect Crit Care*. 2011;71(6):1512–1518.
47. Fantus RJ, Mellett MM, Kirby JP. Use of controlled fascial tension and an adhesion preventing barrier to achieve delayed primary fascial closure in patients managed with an open abdomen. *Am J Surg*. 2006;192(2): 243–247.
48. Hu P, Uhlich R, Gleason F, Kerby J, Bosarge P. Impact of initial temporary abdominal closure in damage control surgery: a retrospective analysis. *World J Emerg Surg*. 2018;13:43. Sep 15 [cited 2019 Apr 26];13. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6139137/>.

49. Cheatham ML, Demetriades D, Fabian TC, Kaplan MJ, Miles WS, Schreiber MA, Holcomb JB, Bochicchio G, Sarani B, Rotondo MF. Prospective study examining clinical outcomes associated with a negative pressure wound therapy system and Barker's vacuum packing technique. *World J Surg.* 2013;37(9):2018–2030.
50. Cothren CC, Moore EE, Johnson JL, Moore JB, Burch JM. One hundred percent fascial approximation with sequential abdominal closure of the open abdomen. *Am J Surg.* 2006;192(2):238–242.
51. Burlew C, Moore E, Biffl W, Bensard D, Johnson J, Barnett C. One hundred percent fascial approximation can be achieved in the postinjury open abdomen with a sequential closure protocol. *J Trauma Acute Care Surg.* 2012;72(1):235–241.
52. Mayberry J, Burgess E, Goldman R, Pearson T, Brand D, Mullins R. Enterocutaneous fistula and ventral hernia after absorbable mesh prosthesis closure for trauma: the plain truth. *J Trauma Inj Infect Crit Care.* 2004;57(1):157–163.
53. Bradley MJ, DuBose JJ, Scalea TM, et al. Independent predictors of enteric fistula and abdominal sepsis after damage control laparotomy: results from the prospective AAST Open Abdomen Registry. *JAMA Surg.* 2013;148(10):947–954.
54. Burlew CC, Moore EE, Cuschieri J, et al. Sew it up! A Western Trauma Association Multi-Institutional Study of enteric injury management in the postinjury open abdomen. *J Trauma Inj Infect Crit Care.* 2011;70(2):273–277.
55. Bilsel Y, Abci I. The search for ideal hernia repair; mesh materials and types. *Int J Surg.* 2012;10(6):317–321.
56. Candage R, Jones K, Luchette FA, Sinacore JM, Vandevender D, Reed RL. Use of human acellular dermal matrix for hernia repair: friend or foe? *Surgery.* 2008;144(4):703–711.
57. Chopra K, Tadisina KK, Matthews JA, Sabino J, Singh DP, Habre W. Early abdominal closure using component separation in patients with an open abdomen after trauma: a pilot study. *Open Access Surgery.* 2015; [cited 2019 Apr 26]. Available from: <https://www.dovepress.com/early-abdominal-closure-using-component-separation-in-patients-with-an-peer-reviewed-fulltext-article-OAS>.
58. Bee T, Croce M, Magnotti L, Zarzaur B, Maish G, Minard G, Schroepel T, Fabian T. Temporary abdominal closure techniques: a prospective randomized trial comparing polyglactin 910 mesh and vacuum-assisted closure. *J Trauma Inj Infect Crit Care.* 2008;65(2):337–344.